



## Critical role of the CD40 CD40-ligand pathway in regulating mucosal inflammation-driven angiogenesis in inflammatory bowel disease

S Danese, F Scaldaferrri, S Vetrano, T Stefanelli, C Graziani, A Repici, R Ricci, G Straface, A Sgambato, A Malesci, C Fiocchi and S Rutella

*Gut* 2007;56:1248-1256; originally published online 22 Feb 2007;  
doi:10.1136/gut.2006.111989

---

Updated information and services can be found at:

<http://gut.bmj.com/cgi/content/full/56/9/1248>

---

*These include:*

### References

This article cites 38 articles, 22 of which can be accessed free at:

<http://gut.bmj.com/cgi/content/full/56/9/1248#BIBL>

1 online articles that cite this article can be accessed at:

<http://gut.bmj.com/cgi/content/full/56/9/1248#otherarticles>

### Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

---

### Notes

---

To order reprints of this article go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Gut* go to:

<http://journals.bmj.com/subscriptions/>

# Critical role of the CD40–CD40-ligand pathway in regulating mucosal inflammation-driven angiogenesis in inflammatory bowel disease

S Danese, F Scaldaferrri, S Vetrano, T Stefanelli, C Graziani, A Repici, R Ricci, G Straface, A Sgambato, A Malesci, C Focchi, S Rutella

*Gut* 2007;56:1248–1256. doi: 10.1136/gut.2006.111989

See end of article for authors' affiliations

Correspondence to: Silvio Danese, MD, Division of Gastroenterology, Istituto Clinico Humanitas, Via Manzoni 56, 20089 Rozzano, Milan, Italy; sdanese@hotmail.com

Revised 13 February 2007  
Accepted 13 February 2007  
Published Online First  
22 February 2007

**Background and aims:** Angiogenesis is a novel component in inflammatory bowel disease (IBD) pathogenesis. We have previously shown that immune–nonimmune interactions through the CD40–CD40-ligand (CD40L) pathway might sustain gut inflammation, although their effect on regulating inflammation-driven angiogenesis is unknown. The present study evaluated the role of the CD40–CD40L interaction in the promotion of immune-mediated angiogenesis in IBD.

**Methods:** Human nonimmune cells of colonic origin—namely, human intestinal fibroblasts (HIFs) and human intestinal microvascular endothelial cells (HIMECs)—were activated with either soluble CD40L (sCD40L), or CD40<sup>+</sup> D1.1 cells or CD40L-activated lamina propria T (LPT) cells before measuring pro-angiogenic cytokine release. Blocking antibodies to either CD40 or CD40L were used to disrupt the CD40–CD40L interaction. The dextran sodium sulphate (DSS) model of experimental colitis in CD40 and CD40L knockout mice was established to assess whether the CD40–CD40L pathway was implicated in controlling inflammation-driven angiogenesis *in vivo*.

**Results:** Engagement of CD40 on HIFs promoted the release of vascular endothelial growth factor (VEGF), interleukin-8 (IL-8) and hepatocyte growth factor (HGF). LPT cells were potent inducers of pro-angiogenic cytokine secretion by HIFs. Supernatants from sCD40L-activated HIFs induced migration of HIMECs and tubule formation, both of which were inhibited by blocking antibodies to either VEGF, IL-8 or HGF. Both CD40- and CD40L-deficient mice were protected from DSS-induced colitis and displayed a significant impairment of gut inflammation-driven angiogenesis, as assessed by microvascular density.

**Conclusions:** The CD40–CD40L pathway appears to be crucially involved in regulating inflammation-driven angiogenesis, suggesting that strategies aimed at blocking CD40–CD40L interactions might be beneficial in acute and chronic intestinal injury.


Angiogenesis is a complex process whereby novel vessels can be generated from pre-existing ones; it involves endothelial cell (EC) division, degradation of vascular basement membrane and surrounding extracellular matrix, and EC migration.<sup>1</sup> Angiogenesis is tightly regulated by a balance between the expression and function of pro-angiogenic and anti-angiogenic mediators. Vascular endothelial growth factor (VEGF) is recognised as a potent pro-angiogenic cytokine and is secreted by monocytes/macrophages, ECs and a variety of other cell types.<sup>2</sup> Similarly, hepatocyte growth factor (HGF) possesses pro-angiogenic properties and promotes tumour cell motility, invasiveness and proliferation.<sup>3,4</sup> Interleukin (IL)-8, initially identified as a pro-inflammatory cytokine, is significantly upregulated in the colonic mucosa of patients with inflammatory bowel disease (IBD), in direct proportion with the degree of inflammation.<sup>5</sup> In addition to being a chemoattractant for neutrophils, IL-8 exerts effects on nonimmune cells, including human intestinal microvascular ECs (HIMECs), is a potent chemoattractant and promotes *in vitro* tube formation in HIMECs by binding IL-8 cognate receptor CXCR2.<sup>6</sup> Both colonic epithelial cells and HIMECs secrete IL-8 in a highly regulated fashion *in vitro*.<sup>7</sup>

Lymphocytes and monocytes might mediate inflammation-driven angiogenesis, which plays a key pathogenic role in several chronic inflammatory diseases, including rheumatoid arthritis, psoriasis and IBD.<sup>8</sup> In such disorders, leukocytes sustain inflammation by producing pro-angiogenic factors and

stimulating angiogenesis.<sup>9,10</sup> We have previously shown that inflammation-driven angiogenesis actively occurs in both Crohn's disease (CD) and ulcerative colitis (UC), being directly correlated with leukocyte infiltrate. Although many of the pathways that trigger inflammatory angiogenesis are unknown at present, the CD40–CD40-ligand (CD40L) system crucially mediates leukocyte–endothelial interactions in the inflamed intestine.<sup>11,12</sup> In particular, both molecules are highly upregulated in IBD, and are functional in the inflamed mucosa by triggering inflammatory cytokine production, cell-adhesion molecule expression and by fostering mucosal inflammation. To date, the role of the CD40–CD40L pathway in mediating inflammatory angiogenesis in the gut remains to be elucidated.

Lymphocytes and monocytes produce pro-angiogenic factors and stimulate angiogenesis during inflammatory responses.<sup>9,10</sup> Excessive production of angiogenesis factors by activated T cells and/or activated macrophages might occur in delayed-type hypersensitivity and in chronic inflammatory disorders.<sup>13</sup>


**Abbreviations:** APCs, antigen-presenting cells; CD, Crohn's disease; CD40L, CD40 ligand; DAI, disease activity index; DSS, dextran sodium sulphate; ECs, endothelial cells; FBS, foetal bovine serum; FGF, fibroblast growth factor; HGF, hepatocyte growth factor; HIFs, human intestinal fibroblasts; HIMECs, human intestinal microvascular endothelial cells; HUVECs, human umbilical-vein endothelial cells; IBD, inflammatory bowel disease; IL-8, interleukin-8; KO, knockout; LP, lamina propria; LPT, lamina propria T cells; sCD40, soluble CD40; TNF, tumour necrosis factor; VEGF, vascular endothelial growth factor; WT, wild type



**Figure 1** VEGF production by CD40L-stimulated HIFs. Panel A: Human intestinal fibroblasts (HIFs) were activated with soluble CD40 ligand (sCD40L) or were co-cultured either with CD40<sup>+</sup> D1.1 cells or with CD40<sup>-</sup> Jurkat cells for 24 hours, before measuring vascular endothelial growth factor (VEGF) release in culture supernatants. In selected experiments, blocking antibodies (Ab) to either CD40 (M2) or CD40L (M90) were used to interfere with the CD40-CD40L interaction. Results are representative of eight independent experiments performed in duplicate and are expressed as mean and standard deviation. \* $p < 0.01$  compared with VEGF release by HIFs not activated with sCD40L; \*\* $p < 0.001$  compared with VEGF release by HIFs cultured in the absence of CD40<sup>+</sup> D1.1 cells; § $p < 0.001$  compared with VEGF release in the absence of blocking antibodies to either CD40 or CD40L. Panel B: HIFs were co-cultured either with unstimulated lamina propria T (LPT) cells or with CD40L<sup>+</sup>-activated LPT cells. In selected experiments, blocking antibodies to either CD40 (M2) or CD40L (M90) were used to interfere with the CD40-CD40L interaction. \*\* $p < 0.001$  compared with VEGF release by HIFs cultured in the absence of CD40<sup>+</sup> LPT; § $p < 0.001$  compared with VEGF release in the absence of blocking antibodies to either CD40 or CD40L.

Interactions between CD40 and CD40L (or CD154) play pluripotent functions in inflammation, predominantly in the effector phase of the immune response.<sup>11-15</sup> CD40, a 50-kD type-I transmembrane-glycoprotein member of the tumour necrosis factor (TNF) receptor gene family, is expressed by numerous cell types, including professional antigen-presenting cells (APCs), monocytes and ECs.<sup>11-17</sup> CD40L, a 33-kD type-II membrane-protein member of the TNF family, is predominantly expressed by activated CD4<sup>+</sup> T cells and platelets.<sup>18</sup> CD40 triggering mediates several biological responses, including immunoglobulin isotype switching in B cells, expression of B7 family molecules on APCs, adhesion-molecule expression by ECs, and expression of cytokines and chemokines that are characteristic of the effector phases of the immune response.


Insights have recently been provided into the role of CD40-CD40L interactions in the induction of VEGF expression and in



**Figure 2** IL-8 production by CD40L-stimulated HIFs. Panel A: Human intestinal fibroblasts (HIFs) were activated with soluble CD40 ligand (sCD40L) or were co-cultured either with CD40<sup>+</sup> D1.1 cells or with CD40<sup>-</sup> Jurkat cells for 24 hours, before measuring interleukin-8 (IL-8) release in culture supernatants. In selected experiments, blocking antibodies (Ab) to either CD40 (M2) or CD40L (M90) were used to interfere with the CD40-CD40L interaction. Results are representative of eight independent experiments performed in duplicate and are expressed as mean and standard deviation. \* $p < 0.001$  compared with IL-8 release by HIFs not activated with sCD40L; \*\* $p < 0.001$  compared with IL-8 release by HIFs cultured in the absence of CD40<sup>+</sup> D1.1 cells; § $p < 0.001$  compared with IL-8 release in the absence of blocking antibodies to either CD40 or CD40L. Panel B: HIFs were co-cultured either with unstimulated lamina propria T (LPT) cells or with CD40L<sup>+</sup>-activated LPT cells. In selected experiments, blocking antibodies to either CD40 (M2) or CD40L (M90) were used to interfere with the CD40-CD40L interaction. \*\* $p < 0.001$  compared with IL-8 release by HIFs cultured in the absence of CD40<sup>+</sup> LPT cells; § $p < 0.001$  compared with IL-8 release in the absence of blocking antibodies to either CD40 or CD40L.

VEGF-stimulated angiogenesis *in vitro* and *in vivo*.<sup>14-19</sup> Human umbilical-vein endothelial cells (HUVECs) and monocytes, but not CD4<sup>+</sup> T cells, release high amounts of VEGF after CD40 triggering by soluble CD40L (sCD40L).<sup>14</sup> Consistent with these findings, injection of human CD40L into SCID mice bearing human skin transplants translates into a significant enhancement of VEGF and fibroblast growth factor (FGF) expression in skin grafts.<sup>19</sup> Finally, blockade of CD40L-CD40 interactions prevents acute and chronic inflammation, including allograft rejection and atherosclerosis.<sup>20-21</sup>

CD40 and CD40L are overexpressed in IBD, suggesting a possible involvement of the CD40-CD40L pathway in disease pathogenesis.<sup>11</sup> Previously, we demonstrated that the CD40-CD40L pathway amplifies intestinal inflammation and links



**Figure 3** HGF production by CD40L-stimulated HIFs. Panel A: Human intestinal fibroblasts (HIFs) were activated with soluble CD40 ligand (sCD40L) or were co-cultured either with CD40<sup>+</sup> D1.1 cells or with CD40<sup>-</sup> Jurkat cells for 24 hours, before measuring hepatocyte growth factor (HGF) release in culture supernatants. In selected experiments, blocking antibodies to either CD40 (M2) or CD40L (M90) were used to interfere with the CD40–CD40L interaction. Results are representative of eight independent experiments performed in duplicate and are expressed as mean and standard deviation. \* $p < 0.001$  compared with HGF release by HIFs not activated with sCD40L; \*\* $p < 0.001$  compared with HGF release by HIFs cultured in the absence of CD40<sup>+</sup> D1.1 cells; § $p < 0.001$  compared with HGF release in the absence of blocking antibodies to either CD40 or CD40L. Panel B: HIFs were co-cultured either with unstimulated lamina propria T (LPT) cells or with CD40L<sup>+</sup> activated LPT cells. In selected experiments, blocking antibodies to either CD40 (M2) or CD40L (M90) were used to interfere with the CD40–CD40L interaction. \*\* $p < 0.001$  compared with HGF release by HIFs cultured in the absence of CD40<sup>+</sup> LPT cells; § $p < 0.001$  compared with HGF release in the absence of blocking antibodies to either CD40 or CD40L.

activated CD40L-expressing T cells with nonimmune cells that, in turn, upregulate cell-adhesion molecules and secrete soluble mediators that recruit T cells, thus sustaining a cycle of immune–nonimmune interactions.<sup>12</sup> Accordingly, disruption of the CD40–CD40L pathway by antibodies to CD40L might favourably affect disease course, as suggested by studies in a murine model of hapten-induced colitis and by studies in patients with IBD.<sup>22–25</sup> It is demonstrated here that CD40 engagement on human nonimmune cells—namely, human intestinal fibroblasts (HIFs) and HIMECs—translates into the promotion of angiogenesis. Furthermore, we provide evidence

that the CD40–CD40L pathway might be involved in the maintenance of inflammation-driven angiogenesis in experimental colitis.


## MATERIALS AND METHODS

### Isolation and culture of HIFs and HIMECs


Surgical specimens used to isolate HIFs and HIMECs were of colonic origin, as reported elsewhere.<sup>12</sup> HIFs were obtained as explants of surgically resected intestinal mucosa, grown to subconfluence in Dulbecco's minimal essential medium supplemented with 10% foetal bovine serum (FBS) and antibiotics, and then established as long-term cultures that were fed twice a week and split at confluence.<sup>12</sup> Isolation of HIMECs consisted of enzymatic digestion of intestinal mucosal strips, followed by gentle compression to extrude EC clumps, which adhered to fibronectin-coated plates and were subsequently cultured in MCDB131 medium (Sigma Aldrich, St Louis, MO) supplemented with 20% FBS, antibiotics, heparin and EC growth factor. For HIF activation, cells were plated onto 24-well cluster plates; after 24–48 hours, the resulting monolayer was fed with fresh medium alone or containing 500 U/mL IFN- $\gamma$  (R&D Systems, Oxon, UK). Cultures of HIFs and HIMECs were maintained at 37 °C in 5% CO<sub>2</sub>. HIFs and HIMECs were used between passage 3 and 10.<sup>12</sup>

### T-cell isolation and culture

Different T-cell preparations were used, including transformed T-cell lines (Jurkat cells; American Type Culture Collection; Manassas, VA) and lamina propria (LP) T cells (LPTs). LP mononuclear cells were isolated as previously detailed.<sup>12–26</sup> Adherent cells were removed by plating  $30 \times 10^6$  cells in RPMI 1640 (Biowhittaker, Walkersville, MD) with 5% FBS onto tissue culture dishes at 37 °C and 5% CO<sub>2</sub> for 1 hour. Nonadherent cells were collected; they were incubated for 30 minutes at 4 °C with magnetically labelled anti-CD19, CD14 and CD16 antibodies (Miltenyi Biotec, Gladbach, Germany) against B cells, monocytes and neutrophils, respectively. T cells were negatively selected by magnetic activated cell sorting.



**Figure 4** Effects of CD40 engagement on HIMEC chemotaxis. Migration of human intestinal microvascular endothelial cells (HIMECs) was measured after activation with either soluble CD40 ligand (sCD40L) or exogenously added vascular endothelial growth factor (VEGF). Additionally, cell migration was assessed after culturing with supernatants from sCD40L-activated human intestinal fibroblasts (HIFs), either in the absence or in the presence of blocking antibodies to VEGF, interleukin-8 (IL-8) or hepatocyte growth factor (HGF). \* $p < 0.001$  compared with baseline HIMEC migration. Results are representative of six independent experiments performed in duplicate and are expressed as mean and standard deviation. § $p < 0.001$  compared with cultures performed in the absence of blocking antibodies to VEGF, IL-8 or HGF.



**Figure 5** Supernatants of CD40L-activated HIF promote *in vitro* tube formation. Tube formation assays were performed with Matrigel™, as detailed in Materials and Methods. Inverted-phase contrast microscopy was used to assess the formation of endothelial cell (EC)-like tube structures. Either vascular endothelial growth factor (VEGF) or soluble CD40 ligand (sCD40L) were used as positive controls for EC formation. Results are representative of three independent experiments. Five high-power fields per culture condition were examined. Magnification  $\times 40$ . HIFs, human intestinal fibroblasts; HIMECs, human intestinal microvascular endothelial cells.

Flow cytometry analysis showed that purified T cells were  $>97\%$  CD3<sup>+</sup>. To induce CD40L expression, LPTs were activated with polystyrene beads coated with murine monoclonal antibodies to human CD3 (10  $\mu\text{g/ml}$ ) and CD28 (1  $\mu\text{g/ml}$ ; DYNAL, Lake Success, NY), as previously reported.<sup>12</sup>

#### Measurement of pro-angiogenic cytokine and chemokine production by HIFs

Confluent HIFs were either left alone or stimulated with 1  $\mu\text{g/ml}$  of sCD40L, or with irradiated D1.1 cells (CD40L<sup>+</sup> acute T-cell leukaemia) or Jurkat cells (CD40<sup>-</sup> acute T-cell leukaemia; 10:1 ratio of T cells to HIFs) for 24 hours. In other wells, a blocking antibody to CD40L (M90 clone; Immunex, Seattle, WA) was used to hinder the CD40-CD40L interaction.<sup>12, 27</sup> Trimeric sCD40L and antibodies against CD40 (M2 clone) were provided by Immunex.

VEGF, HGF and IL-8 levels in HIF-conditioned medium were measured in triplicate with commercially available ELISAs, following the manufacturer's instructions. The limits of detection were as follows:  $<9$  pg/ml VEGF, 3.5 pg/ml IL-8 and  $<40$  pg/ml HGF.


#### HIMEC chemotaxis assay

Chemotaxis was assessed as previously detailed.<sup>28</sup> Briefly, fluorescence-blocked polycarbonate filters (8  $\mu\text{m}$  pore size; BD Biosciences, Palo Alto, CA) were coated with human fibronectin (10  $\mu\text{g/ml}$ ) for 1 hour at room temperature. By using a transwell system,  $35 \times 10^4$  HIMECs were plated in the upper chamber in MCDB-131 medium (Sigma Aldrich), whereas the lower chamber contained chemotaxis buffer with phorbol myristate acetate (10 ng/ml), basic FGF (bFGF; 50 ng/ml) or VEGF (50 ng/ml) as positive controls or 1 ml of supernatants from sCD40L-activated HIFs. In selected wells, 25  $\mu\text{g/l}$  of

blocking antibodies to VEGF, IL-8 or HGF (R&D Systems) or control antibodies were added to the chemotaxis buffer. After 4 hours, buffer was removed from both chambers, and HIMECs migrated onto the lower surface of the porous membrane were washed twice in phosphate-buffered saline and stained with calcein for 15 min at 37 °C. Duplicates of migrated cells were observed with an inverted fluorescence microscope and counted in six random high-power fields. Quantitative analysis of data was performed by using the Ima Pro Plus software (Media Cybernetics, Silver Spring, MD) connected to an Optronics Color digital camera (Olympus Corporation, Tokyo, Japan).

#### Immunostaining and morphometric analysis of intestinal microvasculature

Immunostaining was performed as previously described.<sup>28</sup> Briefly, paraffin-embedded intestinal sections were cut at 3  $\mu\text{m}$  thickness, deparaffinized then hydrated, blocked for endogenous peroxidase using 3% H<sub>2</sub>O<sub>2</sub>/H<sub>2</sub>O, and subsequently subjected to microwave epitope enhancement using a target retrieval solution (Dako, Carpinteria, CA) at pH 10.0. Incubation with anti-CD31 antibodies (Dako) was performed at 1:200 dilution for 30 min at room temperature. Detection was achieved with a standard streptavidin-biotin system (Dako), and antigen localization was visualized with 3'-3-diamino benzidine (Sigma Aldrich). Computerized morphometric analysis was performed using an international consensus method for quantification of angiogenesis.<sup>29</sup> Briefly, stained colonic sections were scanned at low power to detect the most vascularized area, after which at least five microphotographs of the mucosa at high magnification were taken, defined as the area above the muscularis mucosae (ie, the lamina propria), as well as the submucosa, defined as the area between the muscularis mucosae and muscularis propria. The



**Figure 6** Protection from DSS-induced colitis in CD40KO and CD40LKO mice. Experimental colitis was induced with dextran sodium sulphate (DSS) in either wild-type (WT) mice or CD40 knockout (CD40KO) or CD40L knockout (CD40LKO) mice, in accordance with previously published protocols.<sup>33</sup> The percentage of weight loss (panel A) and disease activity index (DAI; panel B) were monitored as indicators of disease severity. Results are representative of independent experiments (eight animals per group) and are expressed as mean and standard deviation. \* $p < 0.01$  when comparing either CD40KO or CD40LKO mice with their WT counterpart.

number of vessels/field (vascular density) was obtained with an Optronics Color digital camera (Olympus). Quantitative analysis of the data was performed with the Image Pro Plus system (Media Cybernetics).

### In vitro tube formation assay

EC tube formation was assessed using Matrigel<sup>TM</sup>, a solubilised extracellular basement membrane matrix extracted from the Engelbreth-Holm-Swarm mouse sarcoma, as detailed elsewhere.<sup>6</sup> Briefly, multi-well dishes were coated with 250  $\mu$ l of complete medium containing 5 mg/ml Matrigel<sup>TM</sup>, and HIMECs resuspended in complete growth medium were seeded at a density of  $5 \times 10^4$ . Cells were cultured on Matrigel<sup>TM</sup> for 16 hours and inverted phase-contrast microscopy was used to assess formation of endothelial tube-like structures. Five

high-power fields per condition were examined and experiments were performed in duplicate.

### Induction and evaluation of colonic inflammation

CD40 knockout (CD40KO) and CD40L knockout (CD40LKO) mice were kindly provided by Mario P. Colombo (Immunotherapy and Gene Therapy Unit, Istituto Nazionale Tumori, Milan, Italy).<sup>30</sup> For the dextran sodium sulphate (DSS) model of colitis, acute inflammation was induced by administration of 3% DSS (40 kDa molecular mass; ICN Biomedicals, Aurora, OH) in filter-purified drinking water for 8 days (Millipore, Bedford, MA). For induction of chronic colitis, mice received three cycles of 5-day 3% DSS followed by 10-day water, as described previously.<sup>31</sup> Mice were sacrificed 10 days after completion of the last cycle. Grading of intestinal inflammation was measured in a blinded fashion by three readers: no inflammation (0); modest numbers of infiltrating cells in the LP (1); infiltration of mononuclear cells leading to separation of crypts and mild mucosal hyperplasia (2); massive infiltration with inflammatory cells accompanied by disrupted mucosal architecture, loss of goblet cells, and marked mucosal hyperplasia (3); all of the above plus crypt abscesses or ulceration (4), with a histological score from 0 to 15.<sup>32</sup> The disease activity index (DAI) was calculated by scoring 1 point for the appearance of each of the following: ruffled fur, occult foecal blood as determined on a Hemocult Sensa card (Smith Kline Diagnostics, San Jose, CA), rectal prolapse  $< 1$  mm, and soft stool. The mice were scored an additional point for diarrhoea or severe rectal prolapse  $> 1$  mm.

### Statistical analysis


The approximation of data distribution to normality was preliminarily tested with statistics for kurtosis and symmetry. Results were presented as mean and SD. All comparisons were performed with the Student's *t*-test for paired or unpaired determinations or with the analysis of variance, as appropriate. The criterion for statistical significance was defined as  $p \leq .05$ .

## RESULTS

### HIFs release pro-angiogenic cytokines on CD40 engagement

In a preliminary set of experiments, we investigated the ability of human CD40-expressing HIFs to release the pro-angiogenic cytokines VEGF, IL-8 and HGF on CD40 triggering. Baseline expression levels and kinetics of CD40 induction in HIFs were extensively described elsewhere.<sup>28</sup> Both sCD40L and CD40L<sup>+</sup> D1.1 cells induced significant VEGF release by HIFs after 24 hours of culture (fig 1A). Conversely, CD40<sup>-</sup> Jurkat cells failed to upregulate VEGF production by co-cultured HIFs, and VEGF levels were comparable to those measured in unstimulated cultures. Blocking antibodies to either CD40L (M90) or CD40 (M2) abrogated VEGF production in response to co-culture with CD40L<sup>+</sup> D1.1 cells, suggesting the involvement of the CD40-CD40L pathway in the promotion of VEGF release in this system model (fig 1A). Interestingly, activated, CD40L-expressing LPT cells, but not resting LPT cells, from healthy subjects promoted VEGF production by HIFs, a response that was significantly inhibited by the addition of antibodies to CD40L or CD40 to the co-cultures (figure 1B). Control antibodies failed to inhibit VEGF production by HIFs (data not shown).

As shown in figure 2A, IL-8 production by HIFs also increased in response to sCD40L and, even more so, to CD40L<sup>+</sup> D1.1 cells. It must be noted that IL-8 was present in the supernatants of co-cultures established with CD40<sup>-</sup> Jurkat cells, although at significantly lower levels compared with cultures supplemented with sCD40L (fig 2A). The addition of



**Figure 7** Protection from histological inflammation after colitis induction in CD40KO and CD40LKO mice. Colonic sections were established from either wild-type (WT) or CD40 knockout (CD40KO) or CD40L knockout (CD40LKO) mice, after development of dextran sodium sulphate (DSS)-induced colitis. Histological damage scores were measured as detailed in Materials and Methods. Results are representative of independent experiments (eight animals per group) and are expressed as mean and standard deviation. \* $p < 0.01$  when comparing either CD40KO or CD40LKO mice with their WT counterpart. Magnification  $\times 10$ .

antibodies to either CD40L or CD40 to the co-cultures of HIF and D1.1 cells translated into a significant reduction of IL-8, the levels of which approached those measured in the presence of CD40<sup>+</sup> Jurkat cells. Similar results were obtained with activated LPT cells (fig 2B). It should be emphasised that both Jurkat cells (fig 2A) and resting LPT cells (fig 2B) elicited modest IL-8 production, reinforcing the concept that IL-8 production by HIFs appears to be only partly dependent on CD40–CD40L interactions.

Finally, sCD40L-stimulated HIFs released significant amounts of HGF and interference with the CD40–CD40L interaction by antibodies specific to either CD40L or CD40 resulted in abrogation of HGF production (fig 3A). In line with the above results on VEGF and IL-8, activated LPT cells were inducers of HGF release by HIFs, and antibodies to CD40 or CD40L significantly, albeit not completely, inhibited HGF secretion (fig 3B). Unexpectedly, bFGF production by HIFs was unaffected by CD40 engagement (data not shown), indicating that the CD40–CD40L pathway might be involved in controlling the release of a distinctive spectrum of pro-angiogenic factors. Collectively, these results pointed to a crucial role for CD40–CD40L interactions in promoting pro-angiogenic cytokine release.

#### CD40 engagement on HIFs promotes EC migration through an indirect mechanism involving pro-angiogenic cytokine release

We next asked whether pro-angiogenic cytokine production by HIFs on CD40 engagement was functional and promoted *in vitro* and *in vivo* angiogenesis. To this end, we transferred culture supernatants of either resting or sCD40L-activated HIFs onto HIMECs before assessing their migratory behaviour. In control cultures, exogenous VEGF was provided as a cytokine stimulus for optimal HIMEC migration. Soluble CD40L induced migration of HIMECs at levels that were comparable to those measured in response to exogenous VEGF (fig 4). Supernatants of HIFs activated with sCD40L for 24 hours induced significant HIMEC migration, which was in part inhibited by blocking

antibodies to either VEGF, IL-8 or HGF, and was completely abrogated by antibodies directed against CD40L (fig 4). These findings suggest that CD40 engagement is capable of promoting HIMEC migration both directly and through the activation of pro-angiogenic cytokine release by HIFs. This conclusion is further corroborated by *in vitro* tube formation experiments. As shown in figure 5, either sCD40L (panel C) or supernatants from CD40L-activated HIFs (panel D) promoted the formation of tube-like structures to a degree comparable to that induced by exogenously added VEGF (panel B). The provision of blocking antibodies to CD40L to cultures of HIMECs and supernatants from sCD40L-activated HIFs inhibited tube formation, indicating a pivotal role for sCD40L in the described phenomena (data not shown).

#### CD40KO and CD40L KO mice manifest diminished angiogenesis and are protected from intestinal inflammation

We next took advantage of the DSS model of experimental colitis to assess whether the CD40–CD40L system might be implicated in controlling colonic inflammation *in vivo* through effects on angiogenesis. As expected, DSS-treated wild-type (WT) mice developed intestinal inflammation and wasting disease.<sup>33</sup> Measurements of both weight loss and DAI clearly indicated that, beginning on day 6 from colitis induction, both CD40KO and CD40LKO mice were significantly protected from DSS-induced colitis compared with the control group of WT mice (fig 6). These findings were underpinned by the histological evaluation and grading of inflammation-induced tissue damage. As shown in figure 7, colonic tissue sections from CD40KO and CD40LKO mice displayed a diminished infiltration with inflammatory cells, and this translated into a lower histological score compared with WT mice. In addition, studies of microvascular density and tissue staining with antibodies to CD31, a well-recognised EC marker, provided objective evidence of diminished angiogenesis in both CD40KO and CD40LKO mice (fig 8). These findings were also confirmed in the chronic model of DSS-induced colitis. As shown in

figure 9, microvascular density was significantly lower in colonic tissue sections of either CD40KO or CD40LKO mice compared with their WT counterpart.

## DISCUSSION


Angiogenesis is a critical component of several chronic inflammatory disorders, including human IBD. However, the pathways that mediate inflammation-associated angiogenesis are not established in the gut. Both CD40 and CD40L are overexpressed in CD and UC, suggesting an intimate involvement in disease pathogenesis and progression.<sup>11</sup> From a therapeutic standpoint, disruption of the CD40–CD40L pathway with subsequent attenuation of gut inflammation could have important implications for human IBD. Indeed, CD40–CD40L-dependent immune–nonimmune interactions favour an influx of pathogenic T cells in the gut mucosa, thus contributing to chronic inflammation.<sup>12</sup> We have recently shown that IBD mucosal extracts induce HIMEC migration and that this response is mostly dependent on IL-8.<sup>28</sup> In addition, IBD mucosal extracts induce potent angiogenic responses in both the corneal and chorioallantoic membrane assays, indicating that the local microvasculature undergoes an intense process of inflammation-dependent angiogenesis.<sup>28</sup> Ligation of CD40 on human pancreatic cells induces secretion of pro-inflammatory chemokines, including IL-8 and MIP-1 $\alpha$ , indicating a potential role of immune–nonimmune interactions in contributing to pancreatic-islet inflammation or graft loss after transplantation.<sup>34</sup> Finally, increased CD40 expression with subsequent induction of IL-8 release has been reported in inflamed human respiratory epithelial cells, further substantiating the view that nonimmune cells can behave as effector cells of the inflammation process following activation of the CD40–CD40L pathway.<sup>35</sup>

The present study was designed to address the role of the CD40–CD40L pathway in inflammation-induced angiogenesis. CD40 engagement on HIFs by either sCD40L or CD40-expressing D1.1 cells induced the release of prototypical pro-angiogenic factors such as VEGF, IL-8 and HGF, whereas CD40-nonexpressing Jurkat cells failed to elicit measurable cytokine production in culture supernatants. Expression of VEGF


following CD40 ligation has been reported in different experimental models, including human umbilical ECs and monocytes.<sup>14</sup> In this study, CD40-dependent VEGF expression was attributed to transcriptional control mechanisms and translated into marked angiogenesis *in vivo*.<sup>14</sup> Specifically, CD40-induced overexpression of VEGF in ECs has been correlated with the activation of the Ras pathway and its effector molecules, providing a mechanism by which the immune system, represented by CD40–CD40L interactions, can regulate angiogenesis.<sup>36</sup>

In our study, LPT cells isolated from the colonic mucosa of healthy subjects and activated *in vitro* through the T-cell receptor to express CD40L, were potent inducers of pro-angiogenic cytokine production by HIFs, suggesting a potential involvement of T cells from inflamed target tissues in the promotion of inflammation-induced angiogenesis. Accordingly, fibroblast-like synovial cells from rheumatoid arthritis patients have been shown to release VEGF after culturing with CD40L-transfected L cells, further supporting the hypothesis that nonimmune cells from inflamed tissues might represent an important source of pro-angiogenic cytokines in a variety of disease models.<sup>37</sup>

Mechanistically, CD40 occupancy on cells of nonimmune origin, such as HIFs, induced pro-angiogenic factor release, which was significantly reduced by neutralizing antibodies directed against either CD40L or CD40. Furthermore, supernatants of HIF cultures were capable of transferring the angiogenesis-inducing activity of CD40-stimulated HIFs, suggesting that cytokine production by HIFs has functional consequences. Blocking antibodies to either VEGF, IL-8 or HGF significantly inhibited HIMEC migration in response to HIF supernatants, unravelling a potential role for these prototypical angiogenic factors in inflammation-driven angiogenic responses. Thus, these experiments point out that the CD40–CD40L pathway can regulate angiogenesis by different mechanisms. First, ECs might be activated by CD40 triggering and this could be the case in CD and UC, in which the elevated levels of sCD40L could induce HIMEC angiogenic activity.<sup>8, 18</sup> The second indirect mechanism might include the stimulation



**Figure 8** Reduction of inflammatory angiogenesis after acute colitis induction in CD40KO and CD40LKO mice. Colonic sections were established from either wild-type (WT) or CD40 knockout (CD40KO) or CD40L knockout (CD40LKO) mice, after development of acute dextran sodium sulphate (DSS)-induced colitis. Microvessel density was used as an indicator of angiogenesis. \* $p < 0.01$  when comparing either CD40KO or CD40LKO mice with their WT counterpart. Magnification  $\times 10$ .



**Figure 9** Reduction of inflammatory angiogenesis after chronic colitis induction in CD40KO and CD40LKO mice. Colonic sections were established from either wild-type (WT) or CD40 knockout (CD40KO) or CD40L knockout (CD40LKO) mice, after development of chronic dextran sodium sulphate (DSS)-induced colitis.<sup>31</sup> Microvessel density was used as an indicator of angiogenesis. \* $p < 0.01$  when comparing either CD40KO ( $n = 8$ ) or CD40LKO ( $n = 7$ ) mice with their WT counterpart ( $n = 7$ ). Magnification  $\times 10$ .

of angiogenic cytokine production by CD40-expressing HIFs, with subsequent activation of HIMECs. Tube-formation experiments substantiated these findings by demonstrating the appearance of tube-like structures after culture of HIMECs in the presence of supernatants from CD40-activated HIFs.

In a final set of experiments, we took advantage of CD40KO and CD40LKO mice to explore the role, if any, of the CD40–CD40L pathway in promoting inflammation-induced angiogenesis *in vivo*. Serial determinations of DAI in CD40KO/CD40LKO mice indicated a significant protection from DSS-induced acute and chronic colonic inflammation when compared with WT animals. Improved intestinal histology and reduced microvascular density in sections of inflamed colons from CD40KO/CD40LKO mice further strengthened the role of CD40–CD40L interactions in inflammation-driven angiogenesis in this IBD experimental model.

In summary, these findings support the notion that immune–nonimmune interactions are important for the maintenance and propagation of inflammation-induced mucosal angiogenesis. We suggest that the modulation of the CD40–CD40L pathway deserves further attention as a future target for therapies to inhibit nonimmune cell-dependent angiogenesis. In mice, administration of CD40L antibodies has translated into a clinical benefit for autoimmune and inflammatory conditions, such as collagen-induced arthritis, lung inflammation and fibrosis, experimental allergic encephalomyelitis, lupus nephritis, atherosclerosis and hapten-induced colitis.<sup>11</sup> Studies with chimeric antagonistic monoclonal antibodies to CD40L (IDEC-131) have been initiated in CD.<sup>25</sup> The administration of IDEC-131 to patients with refractory immune thrombocytopenic purpura has been associated with mild adverse events but not with infections or thromboembolic events,<sup>38</sup> suggesting that this treatment modality might be safe and well tolerated in patients with IBD. Our study also suggests that activated LPT cells bearing CD40L as a consequence of mucosal inflammation might trigger HIF activation and angiogenic cytokine release. Such soluble mediators might, in turn, activate HIMEC angiogenesis. In addition, sCD40L fosters directly mucosal

angiogenesis, pointing to a dual mechanism responsible for CD40-dependent angiogenesis in the inflamed gut. Our studies indicate that therapeutic inhibition of the CD40–CD40L pathway can impact both on nonimmune cell activation and on inflammation-driven angiogenesis in IBD.

## ACKNOWLEDGEMENTS

The authors are indebted to Mario P. Colombo (Immunotherapy and Gene Therapy Unit, Istituto Nazionale Tumori, Milan, Italy) for providing CD40KO and CD40LKO mice.

The authors wish to thank Mr. Renato Zocchi for figure editing.

The authors' studies were supported by grants from the Broad Medical Research Program and by Associazione Italiana per la Ricerca sul Cancro (AIRC) grant (to SD).

## Authors' affiliations

**S Danese, S Vetrano, T Stefanelli, A Repici, A Malesci**, Division of Gastroenterology, Istituto Clinico Humanitas, IRCCS in Gastroenterology, Milan, Italy

**F Scalfaferrì, C Graziani, G Straface**, Department of Internal Medicine, Catholic University Medical School, Rome, Italy

**R Ricci**, Department of Pathology, Catholic University Medical School, Rome, Italy

**A Sgambato**, Institute of General Pathology, Catholic University Medical School, Rome, Italy

**A Malesci**, Division of Gastroenterology, University of Milan, Milan, Italy

**C Fiocchi**, The Cleveland Clinic Foundation, Lerner Research Institute, Cleveland, OH, USA

**S Rutella**, Department of Haematology, Catholic University Medical School, Rome, Italy

Competing interest statement: None declared.

## REFERENCES

- 1 Folkman J. Seminars in Medicine of the Beth Israel Hospital, Boston. Clinical applications of research on angiogenesis. *N Engl J Med* 1995;**333**:1757–63.
- 2 Leung DW, Cachianes G, Kuang WJ, *et al*. Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 1989;**246**:1306–9.
- 3 Rutella S, Bonanno G, Procoli A, *et al*. Hepatocyte growth factor favors monocyte differentiation into regulatory interleukin (IL)-10<sup>+</sup>IL-12<sup>low/neg</sup> accessory cells with dendritic-cell features. *Blood* 2006;**108**:218–27.

- 4 **Jiang WG**, Martin TA, Parr C, *et al*. Hepatocyte growth factor, its receptor, and their potential value in cancer therapies. *Crit Rev Oncol Hematol* 2005;**53**:35–69.
- 5 **Mazzucchelli L**, Hauser C, Zraggen K, *et al*. Expression of interleukin-8 gene in inflammatory bowel disease is related to the histological grade of active inflammation. *Am J Pathol* 1994;**144**:997–1007.
- 6 **Heidemann J**, Ogawa H, Dwinell MB, *et al*. Angiogenic effects of interleukin 8 (CXCL8) in human intestinal microvascular endothelial cells are mediated by CXCR2. *J Biol Chem* 2003;**278**:8508–15.
- 7 **Utgaard JO**, Jahnsen FL, Bakka A, *et al*. Rapid secretion of prestored interleukin 8 from Weibel-Palade bodies of microvascular endothelial cells. *J Exp Med* 1998;**188**:1751–6.
- 8 **Koutroubakis IE**, Tsiolakidou G, Karmiris K, *et al*. Role of angiogenesis in inflammatory bowel disease. *Inflamm Bowel Dis* 2006;**12**:515–23.
- 9 **Polverini PJ**, Cotran PS, Gimbrone MA Jr, *et al*. Activated macrophages induce vascular proliferation. *Nature* 1977;**269**:804–6.
- 10 **Auerbach R**, Sidky YA. Nature of the stimulus leading to lymphocyte-induced angiogenesis. *J Immunol* 1979;**123**:751–4.
- 11 **Danese S**, Sans M, Fiocchi C. The CD40/CD40L costimulatory pathway in inflammatory bowel disease. *Gut* 2004;**53**:1035–43.
- 12 **Vogel JD**, West GA, Danese S, *et al*. CD40-mediated immune-nonimmune cell interactions induce mucosal fibroblast chemokines leading to T-cell transmigration. *Gastroenterology* 2004;**126**:63–80.
- 13 **Ferrara N**. VEGF: an update on biological and therapeutic aspects. *Curr Opin Biotechnol* 2000;**11**:617–24.
- 14 **Melter M**, Reinders ME, Sho M, *et al*. Ligation of CD40 induces the expression of vascular endothelial growth factor by endothelial cells and monocytes and promotes angiogenesis in vivo. *Blood* 2000;**96**:3801–8.
- 15 **Danese S**, Fiocchi C. Platelet activation and the CD40/CD40 ligand pathway: mechanisms and implications for human disease. *Crit Rev Immunol* 2005;**25**:103–21.
- 16 **Rutella S**, Danese S, Leone G. Tolerogenic dendritic cells: Cytokine modulation comes of age. *Blood* 2006;**108**:1435–40.
- 17 **Lederman S**, Yellin MJ, Krichevsky A, *et al*. Identification of a novel surface protein on activated CD4<sup>+</sup> T cells that induces contact-dependent B cell differentiation (help). *J Exp Med* 1992;**175**:1091–101.
- 18 **Danese S**, Katz JA, Saibeni S, *et al*. Activated platelets are the source of elevated levels of soluble CD40 ligand in the circulation of inflammatory bowel disease patients. *Gut* 2003;**52**:1435–41.
- 19 **Reinders ME**, Sho M, Robertson SW, *et al*. Proangiogenic function of CD40 ligand-CD40 interactions. *J Immunol* 2003;**171**:1534–41.
- 20 **Larsen CP**, Elwood ET, Alexander DZ, *et al*. Long-term acceptance of skin and cardiac allografts after blocking CD40 and CD28 pathways. *Nature* 1996;**381**:434–8.
- 21 **Schonbeck U**, Sukhova GK, Shimizu K, *et al*. Inhibition of CD40 signaling limits evolution of established atherosclerosis in mice. *Proc Natl Acad Sci USA* 2000;**97**:7458–63.
- 22 **Stuber E**, Strober W, Neurath M. Blocking the CD40L-CD40 interaction in vivo specifically prevents the priming of T helper 1 cells through the inhibition of interleukin 12 secretion. *J Exp Med* 1996;**183**:693–8.
- 23 **Danese S**, Sans M, Gasbarrini A. CD40 antisense based strategy for inflammatory bowel disease: shutting down multiple cellular communication systems. *Gut* 2005;**54**:891–2.
- 24 **Maerten P**, Liu Z, Ceuppens JL. Targeting of costimulatory molecules as a therapeutic approach in inflammatory bowel disease. *BioDrugs* 2003;**17**:395–411.
- 25 **Dumont FJ**. IDEC-131. IDEC/Eisai. *Curr Opin Investig Drugs* 2002;**3**:725–34.
- 26 **Fiocchi C**, Battisto JR, Farmer RG. Gut mucosal lymphocytes in inflammatory bowel disease: isolation and preliminary functional characterization. *Dig Dis Sci* 1979;**24**:705–17.
- 27 **Danese S**, de la Motte C, Reyes BM, *et al*. Cutting edge: T cells trigger CD40-dependent platelet activation and granular RANTES release: a novel pathway for immune response amplification. *J Immunol* 2004;**172**:2011–5.
- 28 **Danese S**, Sans M, de la Motte C, *et al*. Angiogenesis as a novel component of inflammatory bowel disease pathogenesis. *Gastroenterology* 2006;**130**:2060–73.
- 29 **Vermeulen PB**, Gasparini G, Fox SB, *et al*. Quantification of angiogenesis in solid human tumours: an international consensus on the methodology and criteria of evaluation. *Eur J Cancer* 1996;**32A**:2474–84.
- 30 **Guiducci C**, Valzasina B, Dislich H, *et al*. CD40/CD40L interaction regulates CD4<sup>+</sup>CD25<sup>+</sup> Treg homeostasis through dendritic cell-produced IL-2. *Eur J Immunol* 2005;**35**:557–67.
- 31 **Obermeier F**, Dunger N, Strauch UG, *et al*. CpG motifs of bacterial DNA essentially contribute to the perpetuation of chronic intestinal inflammation. *Gastroenterology* 2005;**129**:913–27.
- 32 **Spencer DM**, Veldman GM, Banerjee S, *et al*. Distinct inflammatory mechanisms mediate early versus late colitis in mice. *Gastroenterology* 2002;**122**:94–105.
- 33 **Kriegelstein CF**, Cerwinka WH, Sprague AG, *et al*. Collagen-binding integrin alpha1beta1 regulates intestinal inflammation in experimental colitis. *J Clin Invest* 2002;**110**:1773–82.
- 34 **Barbe-Tuana FM**, Klein D, Ichii H, *et al*. CD40-CD40 ligand interaction activates proinflammatory pathways in pancreatic islets. *Diabetes* 2006;**55**:2437–45.
- 35 **Cagnoni F**, Oddera S, Giron-Michel J, *et al*. CD40 on adult human airway epithelial cells: expression and proinflammatory effects. *J Immunol* 2004;**172**:3205–14.
- 36 **Flaxenburg JA**, Melter M, Lapchak PH, *et al*. The CD40-induced signaling pathway in endothelial cells resulting in the overexpression of vascular endothelial growth factor involves Ras and phosphatidylinositol 3-kinase. *J Immunol* 2004;**172**:7503–9.
- 37 **Cho CS**, Cho ML, Min SY, *et al*. CD40 engagement on synovial fibroblast up-regulates production of vascular endothelial growth factor. *J Immunol* 2000;**164**:5055–61.
- 38 **Kuwana M**, Nomura S, Fujimura K, *et al*. Effect of a single injection of humanized anti-CD154 monoclonal antibody on the platelet-specific autoimmune response in patients with immune thrombocytopenic purpura. *Blood* 2004;**103**:1229–36.

## EDITOR'S QUIZ: GI SNAPSHOT .....

### Answer

From question on page 1247

The findings in this case are typical of ovarian endometrioma and suspicious for endometriosis of the bowel wall. Various symptoms, such as dysmenorrhea, dyspareunia, chronic pelvic pain, diarrhoea, constipation, cyclic rectal bleeding, colic abdominal pain up to intestinal occlusion may characterise this pathology.

The value of imaging in the diagnosis of endometriosis is limited. Small implants are usually not detectable with ultrasound and are also difficult to identify with MRI. Endometrioma is the only form of endometriosis readily diagnosed with ultrasound (generally transvaginal).

The diagnosis of intestinal endometriosis can be difficult to establish. Endometrial deposits do not invade the mucosa, therefore colonoscopy with biopsies is frequently non-diagnostic. In this case, intestinal ultrasound allowed an endometrial implant in the sigmoid colon to be identified.

It should be noted that endometriosis of the bowel most often involves the sigmoid and rectosigmoid, even though it is a relatively uncommon cause of obstruction. In the event of rectosigmoid strictures of unknown aetiology in premenopausal women, endometriosis should be included in the differential diagnosis. In this case, surgery, which confirmed the diagnosis, was the treatment of choice.

doi: 10.1136/gut.2006.106583a